


Figure 1: A/ Map of plasmid pRIT14586

B/ Coding sequence of the first 127 amino acids
of protein D and multiple cloning site. The signal
sequence is underlined.

BamHI
ATG GAT CCA AAA ACT TTA GCC CTT TCT TTA TTA GCA GCT GGC GTA CTA GCA GGT TGT AGC AGC
 Met Asp Pro Lys Thr Leu Ala Leu Ser Leu Leu Ala Ala Gly Val Leu Ala Gly Cys Ser Ser
 CAT TCA TCA AAT ATG GCG AAT ACC CAA ATG AAA TCA GAC AAA ATC ATT ATT GCT CAC CGT GGT
 His Ser Ser Asn Met Ala Asn Thr Gln Met Lys Ser Asp Lys Ile Ile Ile Ala His Arg Gly
 GCT AGC GGT TAT TTA CCA GAG CAT ACG TTA GAA TCT AAA GCA CTT GCT TTT GCA CAA CAG GCT
 Ala Ser Gly Tyr Leu Pro Glu His Thr Leu Glu Ser Lys Ala Leu Ala Phe Ala Gln Gln Ala
 GAT TAT TTA GAG CAA GAT TTA GCA ATG ACT AAG GAT GGT CGT TTA GTG GTT ATT CAC GAT CAC
 Asp Tyr Leu Glu Gln Asp Leu Ala Met Thr Lys Asp Gly Arg Leu Val Val Ile His Asp His
 TTT TTA GAT GGC TTG ACT GAT GTT GCG AAA AAA TTC CCA CAT CGT CAT CGT AAA GAT GGC CGT
 Phe Leu Asp Gly Leu Thr Asp Val Ala Lys Lys Phe Pro His Arg His Arg Lys Asp Gly Arg
 TAC TAT GTC ATC GAC TTT ACC TTA AAA GAA ATT GAA AGT TTA GAA ATG ACA GAA AAC TTT GAA
 Tyr Tyr Val Ile Asp Phe Thr Leu Lys Glu Ile Gln Ser Leu Glu Met Thr Glu Asn Phe Glu
NcoI ACC ATG GCC ACG TGT GAT CAG AGC TCA SpeI ACT AGT GGA CAC CAT CAC CAT CAC CAT TAA TCT XbaI AGA
 Thr Met Ala Thr Cys Asp Gln Ser Ser Thr Ser Gly His His His His His His *

The amino acid sequence of Figure 1 relates to Seq. ID no. 7 and the nucleic acid sequence of
Figure 1 relates to Seq. ID. No. 6.

The DNA and amino acid sequences of Nef-His; Tat-His; Nef-Tat-His fusion and mutated Tat is illustrated.

Pichia-expressed constructs (plain constructs)

⇒ Nef - HIS

DNA sequence (Seq. ID. No. 8)

ATGGGTGGCAAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGA
ATGAGACGAGCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAA
AAACATGGAGCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGG
CTAGAAGCACAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTA
AGACCAATGACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTAAAAGAAAAGGGG
GGACTGGAAGGGCTAATTCACTCCCAACGAAGACAAGATATCCTTGATCTGTGGATC
TACCACACACAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTC
AGATATCCACTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAG
GTAGAAGAGGCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCAT
GGAATGGATGACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCTAGCA
TTTCATCACGTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGGC
CACCATCACCATCACCATTAA

Protein sequence (Seq. ID. No. 9)

MGGKWSKSSVVGWPTVRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAW
LEAQEEEEVGFPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDLWI
YHTQGYFPDWQNYTPGPGVRYPLTFGWCYKLVPVEPDKVEEANKGENTSLHPVSLH
GMDDPEREVLEWRFD SRLAFHHVARELHPEYFKNCTSGHHHHHH.

⇒ Tat - HIS

DNA sequence (Seq. ID. No. 10)

ATGGAGCCAGTAGATCCTAGACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAA
ACTGCTTGTAACCAATTGCTATTGTAAAAAGTGTTGCTTTTCATTGCCAAGTTTGTTC
ATAACAAAAGCCTTAGGCATCTCCTATGGCAGGAAGAAGCGGAGACAGCGACGAAGA
CCTCCTCAAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAA

TCCCGAGGGGACCCGACAGGCCCGAAGGAACTAGTGGCCACCATCACCATCACCAT
TAA

Protein sequence (Seq. ID. No. 11)

MEPVDPRLEPWKHPGSQPKTACTNCYCKKCCFHCQVCFITKALGISYGRKKRRQRRR
PPQGSQTHQVSLSKQPTSQSRGDPTGPKETSGHHHHHH.

⇒ Nef - Tat - HIS

DNA sequence (Seq. ID. No. 12)

ATGGGTGGCAAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGA
ATGAGACGAGCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAA
AAACATGGAGCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGG
CTAGAAGCACAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTA
AGACCAATGACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTTAAAAGAAAAGGGG
GGACTGGAAGGGCTAATTCCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATC
TACCACACACAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTC
AGATATCCACTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAG
GTAGAAGAGGCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCAT
GGAATGGATGACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCTAGCA
TTTCATCACGTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGAG
CCAGTAGATCCTAGACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAACTGCT
TGTACCAATTGCTATTGTAAAAAGTGTTGCTTTTCATTGCCAAGTTTGTTCATAACA
AAAGCCTTAGGCATCTCCTATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCT
CAAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCCGA
GGGGACCCGACAGGCCCGAAGGAACTAGTGGCCACCATCACCATCACCATTAA

Protein sequence (Seq. ID. No. 13)

^^
MGGKWSKSSVVGWPTVRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAW
LEAQEEEEVGFPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWI
YHTQGYFPDWQNYTPGPGVRYPLTFGWCYKLVPEPDKVEEANKGENTSLLHPVSLH
GMDDPEREVLEWRFD SRLAFHHVARELHPEYFKNCTSEPVDPRLEPWKHPGSQPKTA
CTNCYCKKCCFHCQVCFITKALGISYGRKKRRQRRRPPQGSQTHQVSLSKQPTSQSR
GDPTGPKETSGHHHHHH.

E.coli-expressed constructs (fusion constructs)

⇒ LipoD-Nef-HIS

DNA sequence (Seq. ID. No. 14)

Nucleotides corresponding to the Prot D Fusion Partner are in bold.

The Lipidation Signal Sequence is underlined. After processing, the cysteine coded by the TGT codon, indicated with a star, becomes the amino terminal residue which is then modified by covalently bound fatty acids.

*

ATGGATCCAAAACTTTAGCCCTTTCTTTATTAGCAGCTGGCGTACTAGCAGGTTGT
 AGCAGCCATTTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAAATCATTATT
 GCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATACGTTAGAATCTAAAGCACTT
 GCTTTTGCACAACAGGCTGATTATTTAGAGCAAGATTAGCAATGACTAAGGATGGT
 CGTTTAGTGGTTATTCACGATCACTTTTATAGATGGCTTGACTGATGTTGCGAAAAAA
 TTCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTTACCTTAAAA
 GAAATTCAAAGTTTAGAAATGACAGAAAACCTTTGAAACCATGGGTGGCAAGTGGTCA
 AAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGAGCTGAGCCA
 GCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGAGCAATCACA
 AGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCACAAAGAGGAG
 GAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATGACTTACAAG
 GCAGCTGTAGATCTTAGCCACTTTTTTAAAGAAAAGGGGGGACTGGAAGGGCTAATT
 CACTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACACAAGGCTAC
 TTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTGAGATATCCACTGACCTTT
 GGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAA
 GGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGATGACCCTGAG
 AGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTTCATCACGTGGCCCGA
 GAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGGCCACCATCACCATCACCAT
 TAA

Protein sequence of the processed lipidated ProtD-Nef-HIS protein (Seq. ID. No. 15)

(Amino-acids corresponding to Prot D fusion partner are in bold)

CSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYLEQDLAMTRD
 GRLVVIHDHFLDGLTDVAKKFPHRHRKDGRYYVIDFTLKEIQSLEMTENFETMGGKW
 SKSSVVGWPTVRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAWLEAQE
 EEEVGFPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWYHTQG
 YFPDWQNYTPGPGVRYPLTFGWICYKLVPEPDKVEEANKGENTSLLHPVSLHGMDDP
 EREVLEWRFD SRLAFHHVARELHPEYFKNCTSGHHHHHH.

⇒ LipoD-Nef-Tat-HISDNA sequence (Seq. ID. No. 16)

Nucleotides corresponding to the Prot D Fusion Partner are in bold.

The Lipidation Signal Sequence is underlined. After processing, the cysteine coded by the TGT codon, indicated with a star, becomes the amino terminal residue which is then modified by covalently bound fatty acids.

*

ATGGATCCAAAACTTTAGCCCTTTCTTTATTAGCAGCTGGCGTACTAGCAGGTTGT
 AGCAGCCATTTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAAATCATTATT
 GCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATACTGTTAGAATCTAAAGCACTT
 GCGTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACTAAGGATGGT
 CGTTTAGTGGTTATTCACGATCACTTTTAGATGGCTTGACTGATGTTGCGAAAAAA
 TTCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTTACCTTAAAA
 GAAATTCAAAGTTTAGAAATGACAGAAAACCTTTGAAACCATGGGTGGCAAGTGGTCA
 AAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGAGCTGAGCCA
 GCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGAGCAATCACA
 AGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCACAAAGAGGAG
 GAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATGACTTACAAG
 GCAGCTGTAGATCTTAGCCACTTTTTTAAAAGAAAAGGGGGGACTGGAAGGGCTAATT
 CACTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACACAAGGCTAC
 TTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTGAGATATCCACTGACCTTT
 GGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAA
 GGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGATGACCCTGAG
 AGAGAAGTGTTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTTCATCACGTGGCCCGA
 GAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGAGCCAGTAGATCCTAGACTA
 GAGCCCTGGAAGCATCCAGGAAGTCAGCCCTAAAACCTGCTTGTACCAATTGCTATTGT
 AAAAAGTGTTGCTTTCATTGCCAAGTTTGTTCATAACAAAAGCCTTAGGCATCTCC
 TATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCTCAAGGCAGTCAGACTCAT
 CAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCCGAGGGGACCCGACAGGCCCG
 AAGGAACTAGTGGCCACCATCACCATCACCATTAA

Protein sequence of the processed lipidated ProtD-NEF-TAT-HIS protein (Seq. ID. No. 17)

(Amino-acids corresponding to Prot D fusion partner are in bold)

CSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYLEQDLAMTKD
 GRLVVIHDHFLDGLTDVAKKFPHRHRKDGRIYVIDFTLKEIQSLEMTENFETMGGKW
 SKSSVVGWPTVRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAWLEAQE
 EEEVGFPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWYIYHTQG
 YFPDWQNYTPGPGVRYPLTFGWCYKLPVEPDKVEEANKGENTSLLHPVSLHGMDDP
 EREVLEWRFD SRLAFHHVARELHPEYFKNCTSEPVDPRLEPWKHPGSQPKTACTNCY
 CKKCCFHCQVCFITKALGISYGRKKRRQRRRPPQGSQTHQVSLSKQPTSQSRGDP
 TGPKETSGHHHHHH.

6/17

⇒ ProtD-Nef-HIS

DNA sequence (Seq. ID. No. 18)

Nucleotides corresponding to the Prot D Fusion Partner are in bold.

ATGGATCCAAGCAGCCATTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAA
ATCATTATTGCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATACGTTAGAATCT
AAAGCACTTGCCTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACT
AAGGATGGTCGTTTAGTGTTATTCACGATCACTTTTTAGATGGCTTGACTGATGTT
GCGAAAAAATTCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTT
ACCTTAAAAGAAATTCAAAGTTTAGAAATGACAGAAAACTTTGAAACCATGGGTGGC
AAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAAGAATGAGACGA
GCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGA
GCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCA
CAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATG
ACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTAAAGAAAAGGGGGGACTGGAA
GGGCTAATTCCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACA
CAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTGAGATATCCA
CTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAG
GCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGAT
GACCCTGAGAGAGAAGTGTTAGAGTGAGGTTTGACAGCCGCCTAGCATTTTCATCAC
GTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGGCCACCATCAC
CATCACCATTAA

Protein sequence (Seq. ID. No. 19)

(Amino-acids corresponding to Prot D fusion partner are in bold)

MDPSSHSSNMANTQMKSDKIIAHRGASGYLPEHTLESKALAFQAQADYL
EQDLAMTKDGRLLVVIHDHFLDGLTDVAKKFPHRHRKDGRIYVIDFTLK
EIQSLEMENFETMGGKWSKSSVVGWPTVRERMRRRAEPAADGVGAASRDL
EKHGAITSSNTAATNAACAWLEAQEEEEVGFPVTPQVPLRPMTYKAAVDLSH
FLKEKGGLEGLIHSQRRQDILDWYHTQGYFPDWQNYTPGPGVRYPLTFGW
CYKLVPVEPDKVEEANKGENTSLLHPVSLHGMDDPEREVLEWRFDLSRLAFH
HVARELHPEYFKNCTSGHHHHHH.

⇒ ProtD-Nef-Tat-HIS

DNA sequence (Seq. ID. No. 20)

7/17

Nucleotides corresponding to the Prot D Fusion Partner are in bold.

ATGGATCCAAGCAGCCATTCATCAAATATGGCGAATACCCAAATGAAATCAGACAAA
ATCATTATTGCTCACCGTGGTGCTAGCGGTTATTTACCAGAGCATACGTTAGAATCT
AAAGCACTTGCGTTTGCACAACAGGCTGATTATTTAGAGCAAGATTTAGCAATGACT
AAGGATGGTCGTTTAGTGGTTATTCACGATCACTTTTTAGATGGCTTGACTGATGTT
GCGAAAAAATCCACATCGTCATCGTAAAGATGGCCGTTACTATGTCATCGACTTT
ACCTTAAAGAAATTCAAAGTTAGAAATGACAGAAAACTTTGAAACCATGGGTGGC
AAGTGGTCAAAAAGTAGTGTGGTTGGATGGCCTACTGTAAGGGAAGAATGAGACGA
GCTGAGCCAGCAGCAGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACATGGA
GCAATCACAAGTAGCAATACAGCAGCTACCAATGCTGCTTGTGCCTGGCTAGAAGCA
CAAGAGGAGGAGGAGGTGGGTTTTCCAGTCACACCTCAGGTACCTTTAAGACCAATG
ACTTACAAGGCAGCTGTAGATCTTAGCCACTTTTTAAAAGAAAAGGGGGGACTGGAA
GGGCTAATTCCTCCCAACGAAGACAAGATATCCTTGATCTGTGGATCTACCACACA
CAAGGCTACTTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTGAGATATCCA
CTGACCTTTGGATGGTGCTACAAGCTAGTACCAGTTGAGCCAGATAAGGTAGAAGAG
GCCAATAAAGGAGAGAACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGGAT
GACCCTGAGAGAGAAGTGTAGAGTGGAGGTTTGACAGCCGCCTAGCATTTCATCAC
GTGGCCCGAGAGCTGCATCCGGAGTACTTCAAGAACTGCACTAGTGAGCCAGTAGAT
CCTAGACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAACTGCTTGTACCAAT
TGCTATTGTAAAAAGTGTTGCTTTTCATTGCCAAGTTTGTTCATAACAAAAGCCTTA
GGCATCTCCTATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCTCAAGGCAGT
CAGACTCATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCCGAGGGGACCCG
ACAGGCCCCGAAGGAACTAGTGGCCACCATCACCATCACCATTAA

Protein sequence (Seq. ID. No. 21)

(Amino-acids corresponding to Prot D fusion partner are in bold)

MDPSSHSSNMANTQMKSDKIIIAHRGASGYLPEHTLESKALAFQAQADYLEQDLAMT
KDGRLLVVIHDHFLDGLTDVAKKFPHRHRKDGRYYVIDFTLKEIQSLEMTENFETMGG
KWSKSSVVGWPTVYRERMRAEPAADGVGAASRDLEKHGAITSSNTAATNAACAWLEA
QEEEEVGFPVTPQVPLRPMTYKAAVDLSHFLKEKGGLEGLIHSQRRQDILDLIYHT
QGYFPDWQNYTPGPGVRYPLTFGWCYKLVPVEPDKVEEANKGENTSLLHPVSLHGMD
DPEREVLEWRFD SRLAFHHVARELHPEYFKNCTSEPVDPRLEPWKHPGSQPKTACTN
CYCKKCCFHCQVCFITKALGISYGRKKRRQRRRPPQGSQTHQVSLSKQPTSQSRGDP
TGPKETSGHHHHHH.

⇒ Tat-MUTANT-HIS

DNA sequence (Seq. ID. No. 22)

```

ATGGAGCCAGTAGATCCTAGACTAGAGCCCTGGAAGCATC 40
CAGGAAGTCAGCCTAAACTGCTTGTACCAATTGCTATTG 80
TAAAAAGTGTTGCTTTTCATTGCCAAGTTTGTTCATAACA 120
GCTGCCTTAGGCATCTCCTATGGCAGGAAGAAGCGGAGAC 160
AGCGACGAAGACCTCCTCAAGGCAGTCAGACTCATCAAGT 200
TTCTCTATCAAAGCAACCCACCTCCCAATCCAAAGGGGAG 240
CCGACAGGCCCCGAAGGAACTAGTGGCCACCATCACCATC 280
ACCATTAA 288

```

Protein sequence(Seq. ID. No. 23)

Mutated amino-acids in Tat sequences are in bold.

```

MEPVDPRLPEWKHPGSQPKTACTNCYCKKCCFHCQVCFIT 40
AALGISYGRKKRRRQRRRPPQGSQTHQVLSKQPTSQSKGE 80
PTGPKETSGHHHHHH. 95

```

⇒Nef-Tat-Mutant-HIS

DNA sequence(Seq. ID. No. 24)

```

ATGGGTGGCAAGTGGTCAAAAAGTAGTGTGGTTGGATGGC 40
CTACTGTAAGGGAAAGAATGAGACGAGCTGAGCCAGCAGC 80
AGATGGGGTGGGAGCAGCATCTCGAGACCTGGAAAAACAT 120
GGAGCAATCACAAGTAGCAATACAGCAGCTACCAATGCTG 160
CTTGTGCCTGGCTAGAAGCACAAGAGGAGGAGGAGGTGGG 200
TTTTCAGTCACACCTCAGGTACCTTTAAGACCAATGACT 240
TACAAGGCAGCTGTAGATCTTAGCCACTTTTAAAAGAAA 280
AGGGGGGACTGGAAGGGCTAATCACTCCCAACGAAGACA 320
AGATATCCTTGATCTGTGGATCTACCACACACAAGGCTAC 360
TTCCCTGATTGGCAGAACTACACACCAGGGCCAGGGGTCA 400
GATATCCACTGAÇCTTTGGATGGTGCTACAAGCTAGTACC 440
AGTTGAGCCAGATAAGGTAGAAGAGGCCAATAAAGGAGAG 480
AACACCAGCTTGTTACACCCTGTGAGCCTGCATGGAATGG 520
ATGACCCTGAGAGAGAAGTGTTAGAGTGGAGGTTTGACAG 560
CCGCCTAGCATTTTCATCACGTGGCCCGAGAGCTGCATCCG 600
GAGTACTTCAAGAACTGCACTAGTGAGCCAGTAGATCCTA 640
GACTAGAGCCCTGGAAGCATCCAGGAAGTCAGCCTAAAC 680
TGCTTGTAACCAATTGCTATTGTAAAAAGTGTGCTTTCAT 720
TGCCAAGTTTGTTTCATAACAGCTGCCTTAGGCATCTCCT 760
ATGGCAGGAAGAAGCGGAGACAGCGACGAAGACCTCCTCA 800
AGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAACCC 840
ACCTCCCAATCCAAAGGGGAGCCGACAGGCCCGAAGGAAA 880
CTAGTGGCCACCATCACCATCACCATTAA 909

```



9/17

Protein sequence (Seq. ID. No. 25)


Mutated amino-acids in Tat sequence are in bold.

MGGKWSKSSVVGWPTVRERMRAEPAADGVGAASRDLEKH 40
GAITSSNTAATNAACAWLEAQEEEEVGFPVTPQVPLRPMT 80
YKAAVDLSHFLKEKGGLEGLIHSQRRQDILDWYHTQGY 120
FPDWQNYTPGPGVRYPLTFGWCYKLVPEPDKVEEANKGE 160
NTSLLHPVSLHGMDDPEREVLEWRFD SRLAFHHVARELHP 200
EYFKNCTSEPVDPRLEPWKHPGSQPKTACTNCYCKKCCFH 240
CQVCFITAALGISYGRKKRRQRRRPPQGSQTHQVSLSKQP 280
TSQSKGEPTGPKETSGHHHHHH. 302


11/17

Fig. 4 SDS-PAGE: Nef-Tat-his fusion protein

Daiichi Silver Staining



12/17

Fig . 5 SDS-PAGE: Nef-Tat-his fusion proteinCoomassie blue G250

1: MW (175/83/62,5/47,5/32,5/25/16,5/6,5 kDa)

2: TNH/23 SP eluate (4 µg)

3: TNH/23 Superdex200 eluate (4 µg)

4: TNH/23 Purified bulk (4 µg)

5: TNH/22 Purified bulk (4 µg)

6: TNH/23 Purified bulk (4 µg) / non reducing conditions

7: TNH/22 Purified bulk (4 µg) / non reducing conditions

Fig. 6A Tat-specific antibody titers and isotypes

group	immunization	midpoint titers					ratio IgG1/IgG2a
		Ig	IgG1	IgG2a	IgG2b		
1	oxydized Tat	353557	135538	98771	98763	1,372	
2	reduced Tat	252275	72087	76273	72014	0,945	
3	oxydized Nef-Tat	246466	179616	60835	53563	2,953	
4	reduced Nef-Tat	91726	73767	30948	20679	2,384	
5	adjuvant only	<4000	<4000	<4000	<4000		




Fig. 6B Tat-specific antibody titers and isotypes

group	immunization	midpoint titers					ratio IgG1/IgG2a
		Ig	IgG1	IgG2a	IgG2b		
1	reduced Tat	212799	123242	62697	55763	1,966	
2	reduced Nef-Tat	75676	84046	18449	11692	4,556	
3	adjuvant only	<4000	<4000	<4000	<4000		




Fig. 7 Antigen-specific lymphoproliferative response of pooled lymph node cells

[3H] Thymidine incorporation in cpm				Data expressed as stimulation index			
		Group 1 reduced Tat	Group 2 reduced Nef-Tat	Group 3 adjuvant only			Group 3 adjuvant only
reduced Tat	5µg/ml	41967	18511	789	reduced Tat	5µg/ml	1
	1µg/ml	37609	32346	415		1µg/ml	1
	0.2µg/ml	27640	23408	397		0.2µg/ml	1
reduced Nef-Tat	5µg/ml	43882	31694	483	reduced Nef-Tat	5µg/ml	1
	1µg/ml	33865	28094	245		1µg/ml	0
	0.2µg/ml	25079	22891	383		0.2µg/ml	1
medium		300	161	571	medium		1






Fig. 8 Cell-binding assay

Fig. 9 Inhibition of cell growth